

Article

Assessment of nutritional composition and heavy metal content in some edible mushroom varieties collected from different areas of Bangladesh

K.M. Mohiuddin*, Md. Mehediul Alam, Md. Taufique Arefin and Istiaq Ahmed

Department of Agricultural Chemistry, Faculty of Agriculture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

*Corresponding author: Dr. K.M. Mohiuddin, Department of Agricultural Chemistry, Faculty of Agriculture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh. Mobile: +880-1818675646; E-mail: mohiagchem@gmail.com

Received: 15 November 2015/Accepted: 13 December 2015/ Published: 30 December 2015

Abstract: Four edible mushroom species (*Pleurotus ostreatus*, *Agaricus bisporus*, *Volvariella volvacea*, *Ganoderma lucidum*) from different locations of Bangladesh, were analysed for their protein and metal content profile (K, Na, Fe, Cu, Zn, Mn, Cr, Pb, As and Cd). Trace metals were determined by atomic absorption spectrophotometer, Na and K by flame emission spectrophotometer and protein by micro Kjeldhal method. All element concentrations were determined on a dry weight basis. The protein content of mushrooms varied from 13.8%–34.3% and the metal content of samples ranged from 0.54–2.25% for K and 12.6–81.6, 69.5–626.2, 39.2–163.4, 30.1–75.5, 52.9–104.5, 0.20–0.30, 0.13–0.59 $\mu\text{g g}^{-1}$ for Na, Fe, Cu, Zn, Mn, Cd, Pb, respectively. Arsenic and cadmium concentrations were below the detection limit of the method used. The detection limits of the method for As and Cd are 0.01 $\mu\text{g g}^{-1}$ for each element. In general, K and Fe content were higher than other metals in all mushroom species. The levels of Cu and Zn in some mushroom samples were found to be higher than legal limits.

Keywords: edible mushrooms; heavy metal; nutritional composition

1. Introduction

Mushrooms are large reproductive structures of edible fungi and have been considered as a special kind of food since earliest time and its cultivation is gaining popularity in Bangladesh in recent times. These are very nutritious food and enriched with protein, vitamins and minerals and poor in calorie and cholesterol (Pathak *et al.*, 1998; Chandha and Sharma, 1995; Chang and Miles, 1988 and Bano and Rajarathnam, 1986). Mushrooms have also been reported as therapeutic foods, useful in preventing diseases such as hypertension, hypercholesterolemia (Turkecul *et al.*, 2004) and cancer (Sesli *et al.*, 2008). These functional characteristics are mainly due to their chemical composition (Manzi *et al.*, 2001). Heavy metals are an important source of food contamination and health hazard. The main threats to human health are associated with exposure to As, Cd, Pb, Hg and Cu. Living organisms require trace amounts of some heavy metals, including Fe, Co, Cu, Mn, Cr and Zn. Excessive levels of these metals, however, can be detrimental to living organisms. Other heavy metals such as Cd and Pb have no known beneficial effect on organisms (Falusi and Olanipekun, 2007; Ouzouni *et al.*, 2009; Tuzen *et al.*, 2007). Mushroom species have been known to accumulate great concentrations of heavy metals such as Pb, Cd, Fe, Cu, Mn, Zn, Cr, Ni, Al, and Hg (Kalac *et al.*, 1991; 2001; Svoboda *et al.*, 2000; Kalac and Svoboda, 2001; Falandysz *et al.*, 2003; Dursun *et al.*, 2006; Cocchia *et al.*, 2006; Chen *et al.*, 2009). Recently, extensive research has been carried out all over the world on the factors responsible for the occurrence of heavy metals in mushrooms (Radulescu *et al.*, 2010, Çayır *et al.*, 2010; Genççelep *et al.*, 2009). The accumulation of heavy metals in macrofungi has been found to be affected by environmental and fungal factors. Environmental factors, such as organic matter amount, pH, and metal concentrations in soil, and fungal factors, such as species of mushroom, morphological part of fruiting body,

development stages, age of mycelium, and biochemical composition, affect metal accumulation in macrofungi (Garcia *et al.*, 1998; Kalac and Svoboda, 2001). Heavy metal contaminated food stuff is a great concern in Bangladesh. Higher content of metals have been observed in mushrooms growing in heavily polluted areas, such as close proximity to high ways with heavy traffic, landfills of sewage sludge and emission areas including cities. High content of metals in mushrooms are also reported from areas contaminated by ores mining and processing (Kalac *et al.*, 2004). Thus, a critical assessment of the health risk from mushroom consumption is necessary through screening of metal accumulating varieties. However, there has been no report, to our knowledge, on the heavy metal levels in edible mushroom samples in Bangladesh. The aim of this study was to document basic chemical and metal concentrations in some edible mushrooms, collected in different districts of Bangladesh.

2. Materials and Methods

The mushroom samples were collected from from National Mushroom Development and Extension Centre, Dhaka and Mushroom Sub-center of Comilla, Chittagong, Mymensingh, Sylhet and Khulna city of Bangladesh during early 2014. The collected samples were washed with deionized water and dried at 105 °C for 24 hrs. Dried samples were homogenized using an agate homogenizer and stored in pre-cleaned polyethylene bottles until the analysis started. All reagents were of analytical reagent grade, 69–72% HNO₃, 30% H₂O₂, and 70% HClO₄ were used for digestion of samples. Double deionized water was used for all dilutions. During the experiments, all glasswares and equipment were carefully cleaned starting with 2% HNO₃ and ending with repeated rinsing distilled deionized water to prevent contamination. Samples (0.6 g) were digested with 10 mL of HNO₃, 2 mL of H₂O₂, and 4 mL of HClO₄ in block digester system maintained temperature 180-200 °C until white fumes are evolved and finally diluted to 50 mL with 2% nitric acid. All sample solutions were clear. A blank digest was carried out in the same way. Heavy metals (Fe, Cu, Zn, Mn, Cr, Pb, As and Cd) concentration in the extract were determined by atomic absorption spectrophotometer (SHIMADZU AA-7000), using a deuterium background correction. The crude protein content of the samples was estimated by the micro Kjeldhal method, in which the sample was digested with a known quantity of concentrated sulphuric acid in the block digester. The digested material was distilled after the addition of alkali. The released ammonia was collected in 4% boric acid in the distillation unit. The resultant boric acid, now contained the ammonia released from the digested material, was then titrated against 0.1 N HCl, manually. The nitrogen content thus determined was multiplied by a factor of 6.25 to arrive at the amount of crude protein. Sodium and potassium were determined by flame emission spectrophotometer.

3. Results and Discussion

The habitat, families and common name of mushroom species are listed in Table 1. All of the analysed mushrooms were identified as edible fungi, belonging to the class Agaricomycetes. Basic chemical composition and trace metal levels in the analysed samples are shown in Table 2, expressed in a dry weight basis. The mean levels of protein and analyte ions in analyzed mushroom species are presented in Figure 1. The relative standard deviations were less than 7% for all elements.

Protein concentrations of the mushroom species were generally high and in the range between 13.8 to 34.3%. Protein concentrations of mushroom samples in the literature have been reported to be in the range between 21.57 to 34.77% (Ouzouni *et al.*, 2009), 24.03 to 34.35% (Shelly *et al.*, 2008). Mushrooms proved to be good sources of protein compared with green vegetables.

Mushrooms are also rich in mineral contents. The mushroom species provides a reasonable amount of minerals in comparison with vegetables (Guillamón *et al.*, 2010). The evaluated strains differed in mineral content. The levels of K in the samples ranged from 0.54 to 2.25%, and the highest K levels were obtained in PHK strain of Oyster mushroom. Potassium concentration has been reported to be 3.10 to 3.7% in the literature (Shelly *et al.*, 2008). Minimum and maximum concentrations of Na were measured as 12.6 and 81.6 µg g⁻¹ in PHK strain and PO2 strain of Oyster mushroom, respectively. Sodium concentrations of mushroom samples, in the literature, have been reported in the ranges of 0.03–4.85 mg g⁻¹ (Genccelep *et al.*, 2009). The concentration of Na is relatively low and this is of very great nutritional benefit to the consumer, a finding that has been corroborated by Vetter (2003).

Table 1. Families, Habitat, edibility and common name of mushroom species.

No.	Species	Family	Habitat	Common name
1	<i>Pleurotus ostreatus</i> (Jacq. ex Fr.) P. Kumm	Pleurotaceae	In mixed woods	Oyster
2	<i>Agaricus bisporus</i> (J.E. Lange) Imbach	Agaricaceae	In forests	Button
3	<i>Volvariella volvacea</i> (Bul. ex Fr.) Singer	<u>Pluteaceae</u>	In forests	Straw
4	<i>Ganoderma lucidum</i> (Curtis) P. Kars	Ganodermataceae	On trees	Reishi

The levels of trace metals Fe, Cu, Zn, Mn, Cr, Pb in mushroom species were found to be 69.5–626.2, 39.2–163.4, 30.1–75.5, 52.9–104.5, 0.20–0.30, 0.13–0.59 $\mu\text{g g}^{-1}$, respectively. Arsenic and cadmium contents were below detectable limit determined by flame AAS method. The order of the mean levels of heavy metals in the mushroom samples was found to be as Fe > Cu > Mn > Zn > Cr > Pb. The FAO/WHO has set a limit for heavy metals intakes based on body weight. For an average adult (60 kg body weight), the provisional tolerable daily intake for Fe, Cu, Zn, and Pb are 48 mg, 3 mg, 60 mg, and 214 $\mu\text{g g}^{-1}$, respectively (FAO/WHO 1999). The trace metal contents in the mushrooms are mainly affected by acidic and organic matter content of their ecosystem and soil (Gast *et al.*, 1988). The uptake of metal ions in mushrooms is in many aspects different from plants. For this reason, the concentration variations of metals depend on mushroom species and their ecosystems (Chojnacka and Falandysz 2007; Kowalewska *et al.*, 2007; Shin *et al.*, 2007).

Iron is vital for almost all living organisms, participating in a wide variety of metabolic processes, including oxygen transport, DNA synthesis, and electron transport (Lynch and Baynes, 1996). The range of Fe levels were between 69.5–626.2 $\mu\text{g g}^{-1}$, most samples having concentrations between 130 and 400 $\mu\text{g g}^{-1}$. The average concentration of Fe was 188.2 $\mu\text{g g}^{-1}$. The maximum Fe level permitted for food is 15 $\mu\text{g g}^{-1}$ according to Turkish Food Codex Anonymous Regulation (2002). Iron levels in all analyzed mushroom samples were found to be higher than legal limits. The reported Fe values for mushroom samples were 31.3–1,190 $\mu\text{g g}^{-1}$ (Sesli and Tüzen, 1999), 30–150 $\mu\text{g g}^{-1}$ (Kalač and Svoboda, 2001), 180–407 $\mu\text{g g}^{-1}$ (Isiloglu *et al.*, 2001), 146–835 $\mu\text{g g}^{-1}$ (Tüzen, 2003), 56.1–7,162 $\mu\text{g g}^{-1}$ (Mendil *et al.*, 2004), 568–3,904 $\mu\text{g g}^{-1}$ (Türkecul *et al.*, 2004), 102–1,580 $\mu\text{g g}^{-1}$ (Soylak *et al.*, 2005), 211–628 $\mu\text{g g}^{-1}$ (Mendil *et al.*, 2005), 110–11,460 $\mu\text{g g}^{-1}$ (Yamaç *et al.*, 2007), and 150–1,741 $\mu\text{g g}^{-1}$ (Sesli *et al.*, 2008), 67.5–843 $\mu\text{g g}^{-1}$ (Zhu *et al.*, 2011), respectively. Our Fe values are similar to those of previous studies.

Copper is the third-most abundant trace element in human body and small amount of Cu is found in the human body (50–120 mg), but it plays a critical role like vitamin in a variety of biochemical processes (Yaman and Akdeniz 2004). Copper forms part of at least 13 different enzymes, and its presence is needed for each if they are to function properly. It is known that Cu may be toxic to both humans and animals when its concentration exceeds the safe limits (Gast *et al.*, 1988). *P. ostreatus* had the lowest copper concentration (39.2 $\mu\text{g g}^{-1}$) whereas *A. bisporus* had the highest (163.4 $\mu\text{g g}^{-1}$). The average Cu level of the samples was 85.88 $\mu\text{g g}^{-1}$. Copper concentrations, accumulated in mushroom species, are usually 100–300 $\mu\text{g g}^{-1}$, which is not considered a health risk (Soylak *et al.*, 2005, Kalac and Svoboda, 2001). These levels are above the WHO permissible limits in foods, which is 40 $\mu\text{g g}^{-1}$ (Bahemuka and Mubofu, 1999). Copper contents of mushroom samples in the literature have been reported to be in the ranges: 4.71–51.0 $\mu\text{g g}^{-1}$ (Tüzen *et al.*, 1998), 10.3–145 $\mu\text{g g}^{-1}$ (Sesli and Tüzen 1999), 12–181 $\mu\text{g g}^{-1}$ (Tüzen *et al.*, 2003), 12–181 $\mu\text{g g}^{-1}$ (Tüzen, 2003), 13.4–50.6 $\mu\text{g g}^{-1}$ (Soylak *et al.*, 2005), 24.1–86.2 $\mu\text{g g}^{-1}$ (Mendil *et al.*, 2005), 10.6–144.2 $\mu\text{g g}^{-1}$ (Yamaç *et al.*, 2007), and 15–73 $\mu\text{g g}^{-1}$ (Sesli *et al.*, 2008), respectively. Our Cu levels were found to be higher than those reported in the literature. The Cu results of our study were in similar with those found in the literature (Demirbas, 2001a; Isildak *et al.*, 2004; Kalac *et al.*, 1996; Svoboda *et al.*, 2000). Our Cu contents were found to be lower than 6.83–31.9 $\mu\text{g g}^{-1}$ (Zhu *et al.*, 2011).

Zinc is an integral component of a wide variety of different enzymes in which it plays catalytic, structural, and regulatory roles. Minimum and maximum concentrations of Zn in our samples were 30.1 and 75.5 $\mu\text{g g}^{-1}$. The mean Zn content of the samples was 43.24 $\mu\text{g g}^{-1}$. The WHO permissible limit of Zn in foods is 60 $\mu\text{g g}^{-1}$ (WHO 1982). The values for Zn in some investigated mushroom samples were above the WHO's values. Zinc concentrations of mushroom samples in the literature have been reported to be in the ranges: 29.3–158 $\mu\text{g g}^{-1}$ (Isiloglu *et al.*, 2001), 33.5–89.5 $\mu\text{g g}^{-1}$ (Tüzen, 2003), 40.3–64.4 $\mu\text{g g}^{-1}$ (Mendil *et al.*, 2004), 45.2–173.8 $\mu\text{g g}^{-1}$ (Soylak *et al.*, 2005), and 43.5–205 $\mu\text{g g}^{-1}$ (Sesli *et al.*, 2008), 42.9–94.3 $\mu\text{g g}^{-1}$ (Zhu *et al.*, 2011), respectively. Our Zn values are in agreement with literature values.

Table 2. Concentrations of protein (%), K (%), Na ($\mu\text{g g}^{-1}$) and trace metals ($\mu\text{g g}^{-1}$) of the mushroom samples analyzed.

Mushroom species	Area	Location	Strain	Protein	K	Na	Fe	Cu	Zn	Mn	Cr	Pb	As & Cd	
<i>P. ostreatus</i>	Dhaka	NMDEC **	PO2	18.5±0.14	1.75±0.31	32.3±2.1	180.8±2.4	77.6±0.89	35.4±0.32	54.3±0.88	0.21±0.02	0.14±0.02	BDL*	
		NMDEC	Florida	16.7±0.20	1.92±0.11	39.7±1.9	163.5±3.1	96.3±1.30	32.6±0.44	64.8±1.20	0.22±0.01	0.15±0.01	BDL	
		NMDEC	HK51	18.5±0.12	1.73±0.10	69.3±2.8	345.2±4.8	102.1±2.6	35.5±0.28	68.4±1.56	0.23±0.01	0.14±0.02	BDL	
		Agora (SQ)	PO2	25.7±0.15	1.79±0.60	32.3±1.3	114.9±4.2	88.7±1.12	44.2±0.35	104.5±1.8	0.23±0.01	0.18±0.02	BDL	
		Agora (DM)	PO2	19.7±0.08	0.77±0.03	32.3±0.9	164.2±4.9	88.5±1.40	36.2±0.23	70.1±0.90	0.23±0.01	0.59±0.03	BDL	
		Agora (MB)	PO2	20.9±0.18	1.16±0.4	22.4±1.1	196.5±3.5	160.5±1.9	45.0±0.49	83.0±0.85	0.24±0.00	0.30±0.01	BDL	
		Meenabazar		21.0±0.22	1.59±0.02	27.4±1.9	170.0±4.2	39.2±0.88	37.8±0.23	88.0±0.97	0.23±0.01	0.24±0.01	BDL	
		Newmarket		21.6±0.31	1.50±0.08	32.3±2.1	137.6±4.2	57.6±0.65	38.8±0.17	76.0±1.02	0.25±0.02	0.21±0.02	BDL	
		Kawranbazar		21.8±0.16	1.56±0.20	37.2±2.2	166.8±3.4	89.2±0.44	48.6±0.24	75.1±1.14	0.26±0.02	0.36±0.01	BDL	
		Nutunbazar		14.8±0.13	1.75±0.08	47.1±2.0	360.0±5.9	87.5±1.20	30.4±0.18	70.9±0.96	0.27±0.01	0.20±0.01	BDL	
	Rowja Brand		24.3±0.11	2.03±0.62	71.8±3.5	162.5±3.4	56.1±0.97	30.6±0.21	70.1±0.89	0.26±0.01	0.16±0.01	BDL		
	Chittagong	NMDESC ***	PO2	19.0±0.14	1.35±0.21	29.8±2.3	161.8±3.2	79.5±0.85	50.2±0.43	67.1±0.85	0.21±0.01	0.14±0.02	BDL	
		NMDESC	PSC	22.7±0.15	1.32±0.02	32.3±2.4	563.8±6.5	84.5±1.20	56.7±0.42	70.6±0.93	0.20±0.00	0.14±0.02	BDL	
		NMDESC	HK51	20.3±0.09	1.54±0.08	47.1±3.2	69.5±1.6	87.1±1.16	52.9±0.34	52.9±1.04	0.22±0.01	0.14±0.03	BDL	
		NMDESC	PHK	25.8±0.16	2.25±0.52	12.6±0.8	369.0±3.2	73.5±1.08	51.9±0.24	73.0±1.09	0.21±0.00	0.14±0.01	BDL	
	Mymensingh	NMDESC		22.8±0.14	1.52±0.03	47.1±1.2	626.2±5.8	74.3±1.15	75.5±0.54	85.5±1.35	0.27±0.02	0.14±0.02	BDL	
		NMDESC		26±0.18	1.53±0.05	74.2±2.6	193.4±2.3	79.5±0.67	32.8±0.26	79.3±1.24	0.24±0.01	0.14±0.02	BDL	
		Ganginarpar		15.8±0.17	1.14±0.02	76.7±3.5	182.0±2.5	96.0±0.86	30.1±0.19	75.8±1.03	0.27±0.01	0.16±0.03	BDL	
		Seshmor		18.5±0.09	0.75±0.02	27.4±2.1	300.0±4.6	86.0±1.35	44.5±0.42	80.8±1.08	0.27±0.01	0.16±0.01	BDL	
		Rohmotpur		14.6±0.10	0.94±0.02	47.1±3.1	270.8±3.9	55.0±1.02	35.5±0.23	75.0±0.95	0.29±0.01	0.16±0.01	BDL	
		Sankipara		13.8±0.12	0.91±0.01	37.2±1.6	392.4±4.2	84.0±2.10	31.4±0.14	89.3±0.90	0.30±0.01	0.16±0.01	BDL	
	Comilla	NMDESC	PO2	16.7±0.08	1.60±0.08	81.6±3.6	129.5±2.4	81.0±1.23	34.3±0.20	83.8±1.06	0.25±0.01	0.14±0.01	BDL	
		NMDESC	Florida	34.3±0.19	1.56±0.05	32.3±2.1	348.5±3.5	134.6±2.8	61.3±0.47	74.9±1.12	0.26±0.02	0.15±0.02	BDL	
	Rajshahi	NMDESC	PO2	22.5±0.27	1.33±0.04	37.2±1.6	151.2±3.1	74.5±0.46	54.9±0.35	83.5±1.29	0.26±0.01	0.14±0.01	BDL	
	<i>A. bisporus</i>	Dhaka	Agora (SQ)		24.7±0.34	1.37±0.05	57.0±2.9	143.6±4.6	163.4±3.9	47.6±0.46	91.1±1.38	0.29±0.01	0.16±0.02	BDL
			Chainese can-1		17.7±0.13	0.54±0.02	52.0±2.7	396.0±5.3	79.7±1.20	39.6±0.38	80.7±1.24	0.30±0.01	0.22±0.02	BDL
Chainese can-2				18.6±0.12	1.04±0.08	37.2±2.1	370.0±4.8	54.6±0.86	36.3±0.23	83.7±1.23	0.23±0.01	0.15±0.01	BDL	
Chainese can-3				19.6±0.09	1.58±0.04	61.9±3.4	146.0±3.2	71.7±0.23	58.0±0.41	56.2±1.34	0.25±0.00	0.17±0.03	BDL	
<i>V. volvacea</i>			18.9±0.14	1.35±0.03	44.6±2.8	322.5±5.2	101.8±2.3	36.5±0.43	78.5±0.97	0.24±0.01	0.25±0.02	BDL		
<i>G. lucidum</i>			22.8±0.20	0.91±0.01	37.2±1.3	303.0±3.6	72.5±1.22	52.2±0.47	64.0±0.92	0.21±0.01	0.13±0.01	BDL		
Average				20.62	1.40	43.89	253.37	85.88	43.24	75.70	0.25	0.19	BDL	

*BDL= Below detectable limit using flame AAS (0.01 $\mu\text{g g}^{-1}$)

**NMDEC: National Mushroom Development and Extension Centre

***NMDEC: National Mushroom Development and Extension Sub-centre

Table 3. Correlations between heavy metal concentrations of the edible mushroom samples.

	Na	Fe	Cu	Zn	Mn	Cr	Pb
Na	1						
Fe	0.593	1					
Cu	0.888	0.157	1				
Zn	-0.999*	-0.551	-0.911	1			
Mn	-0.931	-0.259	-0.995	0.949	1		
Cr	-0.031	-0.823	0.431	-0.020	-0.335	1	
Pb	0.064	0.842	-0.401	-0.013	0.304	-0.999*	1

*Significance at 5% level of probability




Figure 1. Distribution of mean values of protein and analyte ions in mushroom species: (1) *P. ostreatus*; (2) *A. bisporus*; (3) *A. bisporus*; (4) *G. lucidum*

Manganese is present in metalloproteins, such as pyruvate carboxylase, and in the cytoplasmic glial enzyme, glutamine synthetase. In this study, the lowest Mn content was $52.9 \mu\text{g g}^{-1}$, for the HK51 strain of *P. ostreatus* species, whereas the highest Mn content was $104.5 \mu\text{g g}^{-1}$, for the PO2 strain of *P. ostreatus* species. The mean Mn content of the samples was $75.70 \mu\text{g g}^{-1}$. Toxicity limits of Mn for plants are high ($400\text{--}1,000 \mu\text{g g}^{-1}$). Our Mn levels are below toxicity limits. Manganese concentrations of mushroom samples, in the literature, have been reported in the ranges of $7.6\text{--}56.2 \mu\text{g g}^{-1}$ (Demirbas 2001b), $14.5\text{--}63.6 \mu\text{g g}^{-1}$ (Isiloglu *et al.*, 2001), $5.0\text{--}60.0 \mu\text{g g}^{-1}$ (Kalač and Svoboda, 2001), $12.9\text{--}93.3 \mu\text{g g}^{-1}$ (Tüzen, 2003), $7.1\text{--}81.3 \mu\text{g g}^{-1}$ (Isildak *et al.*, 2004), $21.7\text{--}74.3 \mu\text{g g}^{-1}$ (Mendil *et al.*, 2004), $18.1\text{--}103 \mu\text{g g}^{-1}$ (Mendil *et al.*, 2005), $14.2\text{--}69.7 \mu\text{g g}^{-1}$ (Soylak *et al.*, 2005), $13.5\text{--}113 \mu\text{g g}^{-1}$ (Zhu *et al.*, 2011) respectively. Our values for these species are in agreement with those reported earlier.

Chromium plays vital role for the normal metabolism of cholesterol, fat, and glucose. Chromium contents of mushroom samples were found in $0.20\text{--}0.30 \mu\text{g g}^{-1}$. The average Cr content of the samples was $0.25 \mu\text{g g}^{-1}$. These values were well below the FDA recommended daily intake of Cr for foods and feeds, which is $120 \mu\text{g g}^{-1}$ (Haider *et al.*, 2004). Chromium values in mushroom samples have been reported to be in the ranges: $7.0\text{--}11.0 \mu\text{g g}^{-1}$ (Sivrikaya *et al.*, 2002), $0.87\text{--}2.66 \mu\text{g g}^{-1}$ (Tüzen, 2003), $0.16\text{--}4.86 \mu\text{g g}^{-1}$ (Malinowska *et al.*, 2004), $1.2\text{--}4.2 \mu\text{g g}^{-1}$ (Mendil *et al.*, 2004), $0.34\text{--}1.10 \mu\text{g g}^{-1}$ (Soylak *et al.*, 2005), and $1.95\text{--}73.8 \mu\text{g g}^{-1}$ (Yamaç *et al.*, 2007), $10.7\text{--}42.7 \mu\text{g g}^{-1}$ (Zhu *et al.*, 2011), respectively. Chromium contents were found to be lower than those reported earlier. Lead has no beneficial role in human metabolism, producing progressive toxicity. The Pb level ranged from 0.13 to $0.59 \mu\text{g g}^{-1}$ for *G. lucidum* and strain PO2 of *P. ostreatus*. The average Pb content of the samples was $0.19 \mu\text{g g}^{-1}$. These values were below the WHO permissible limit ($2.0 \mu\text{g g}^{-1}$) in plant. Lead contents of mushroom samples in the literature have been reported to be in the ranges: $0.40\text{--}2.80 \mu\text{g g}^{-1}$ (Svoboda *et al.*, 2000), $1.43\text{--}4.17 \mu\text{g g}^{-1}$ (Tüzen, 2003), $0.800\text{--}2.700 \mu\text{g g}^{-1}$ (Türkekul *et al.*, 2004), $0.82\text{--}1.99 \mu\text{g g}^{-1}$ (Soylak *et al.*, 2005), and $0.9\text{--}2.6 \mu\text{g g}^{-1}$ (Sesli *et al.*, 2008), respectively. The Pb results of all mushroom species were in agreement with those found in the literature. The permissible limit of Cd and As in plants, recommended by WHO, is $0.02 \mu\text{g g}^{-1}$ and our values of Cd and As in mushroom samples were below $0.01 \mu\text{g g}^{-1}$ (below detectable limit by flame AAS method). Statistically significant correlation coefficients at 0.05 probability level were established between metal concentrations. The values of correlation coefficients between metal concentrations are given in Table 3. There are good correlations between Mn and Zn ($r = 0.949$), Na and Cu ($r = 0.888$), Fe and Pb ($r = 0.842$), Na and Fe ($r = 0.593$). There are positive correlations of Na and Fe, Na and Cu, Na and Pb, Fe and Cu, Fe and Pb, Cu and Cr, Zn and Mn, Mn and Pb. Negative correlations were found between Na and Zn, Na and Mn, Na and Cr, Fe and Zn, Fe and Mn, Fe and Cr, Cu and Zn, Cu and Mn, Cu and Pb, Zn and Cr, Zn and Pb, Mn and Cr, Cr and Pb.

4. Conclusions

In the present study, the detected levels of Fe, Zn, Mn and Pb were generally in agreement with previously reported. But the Cu contents were higher than those reported earlier and the Cr levels were lower than literature values. In general, the levels of Cu and Zn in some mushroom samples were found to be higher than legal limits. The heavy metal levels of edible mushrooms should be analyzed more often in Bangladesh in order to evaluate the possible danger to human health from them.

Acknowledgements

The authors would like to thank the University Grants Commission of Bangladesh and Bangladesh Agricultural University Research Systems for financing project No.: 2013/42/BAU-UGC.

Conflict of interest

None to declare.

References

- Anonymous, 2002. Regulation of setting maximum levels for certain contaminants in foodstuffs. Official Gazette, Iss: 24908.
- Bahemuka TE and EB Mubofu, 1999. Heavy metals in edible green vegetables grown along the sites of the Sinza and Msimbazi rivers in Dares Salaam, Tanzania. Food Chem., 66: 63–66.
- Bano Z and S Rajarathnam, 1986. Vitamin values of Pleurotus mushrooms. Qualitas Plantarum Plant Foods for Human Nutrition, 36: 11-15.

- Cayır A, M Coskun and M Coskun, 2010. The heavy metal content of wild edible mushroom samples collected in Canakkale Province, Turkey. *Biol. Trace Elem. Res.*, 134: 212–219.
- Chandha KL and SR Sharma, 1995. *Advances in Horticulture. Mushroom*, Malhotra Publication house, New Delhi. pp. 1-33.
- Chang ST and PG Miles, 1988. *Edible Mushroom and their cultivation*. CRC Press, Inc. Boca Raton, Florida U.S.A. pp. 83–88.
- Chen XH, HB Zhou and GZ Qiu, 2009. Analysis of several heavy metals in wild edible mushrooms from regions of China. *Bull. Environ. Contam. Toxicol.*, 83: 280–285.
- Chojnacka A and J Falandysz, 2007. Mineral composition of yellow-cracking bolete (*Xerocomus subtomentosus*) (L.) Quelet. *Bromat Chem. Toksykol*, 40: 337–340.
- Cocchia L, L Vescovia, LE Petrinid and O Petrini, 2006. Heavy metals in edible mushrooms in Italy. *Food Chem.*, 98: 277–284.
- Demirbas A, 2001a. Concentrations of 21 metals in 18 species of mushrooms growing in the East Black Sea region. *Food Chem.*, 75: 453–457.
- Demirbas A, 2001b. Heavy metal bioaccumulation by mushrooms from artificially fortified soils. *Food Chem.*, 74: 293–301.
- Dursun N, MM Ozcan, G Kasik, and C Ozturk, 2006. Mineral contents of 34 species of edible mushrooms growing wild in Turkey. *J. Sci. Food Agric.*, 86: 1087–1094.
- Falandysz J, M Kawano, A Swieczkowski, A Brzostowski and M Dadej, 2003. Total mercury in wild-grown higher mushrooms and underlying soil from Wdzydze Landscape Park, Northern Poland. *Food Chem.*, 81: 21–26.
- Falusi BA and EO Olanipekun, 2007. Bioconcentration factors of heavy metals in tropical crab (*Carcinus* sp.) from River Aponwe, Ado-Ekiti, Nigeria. *J. Appl. Sci. Environ. Manage.*, 11: 51–54.
- FAO/WHO, 1999. Expert Committee on Food Additives, Summary and Conclusions. Fifty-third Meeting, Rome.
- Garcia MA, J Alonso, MI Fernandez and MJ Melgar, 1998. Lead content in edible wild mushrooms in Northwest Spain as indicator of environmental contamination. *Arch. Environ. Contam. Toxicol.*, 34: 330–335.
- Gast CH, E Jansen, J Bierling and L Haanstra, 1988. Heavy metals in mushrooms and their relationship with soil characteristics. *Chemosphere*, 17: 789–799.
- Gençcelep H, Y Uzun, Y Tunçturk and K Demirel, 2009. Determination of mineral contents of wild-grown edible mushrooms. *Food Chem.*, 113: 1033–1036.
- Guillamón E, A Garcia-Lafuente, M Lozano, M D'Arrigo, MA Rostagno, A Villares, and JA Martínez, 2010. Edible mushrooms: role in the prevention of cardiovascular diseases. *Fitoterapia*, 81: 715-723.
- Haider S, V Naithani, J Barthawal, and P Kakkar, 2004. Heavy metal content in some therapeutically important medicinal plants. *Bull. Environ. Contam. Toxicol.*, 72: 119–127.
- Isildak O, I Turkecul, M Elmastas and M Tüzen, 2004. Analysis of heavy metals in some wild-grown edible mushrooms from the middle Black Sea region, Turkey. *Food Chem.*, 86: 547–552.
- Isiloglu M, F Yilmaz and M Merdivan, 2001. Concentrations of trace elements in wild edible mushrooms. *Food Chem.*, 73: 169–175.
- Kalač P, J Burda and I Staskova, 1991. Concentrations of lead, cadmium, mercury and copper in mushrooms in the vicinity of a lead smelter. *Sci. Total Environ.*, 105: 109–119.
- Kalač P and L Svoboda, 2001. A review of trace element concentrations in edible mushrooms. *Food Chem.*, 69: 273–281.
- Kalač P, L Svoboda and Havlíčková, 2004. Contents of cadmium and mercury in edible mushrooms. *J. Appl. Biomed.*, 2: 15–20.
- Kalač P, M Niznanska, D Bevilaqua, and I Staskova, 1996. Concentrations of mercury, copper, cadmium and lead in fruiting bodies of edible mushrooms in the vicinity of a mercury smelter and a copper smelter. *Sci. Total Environ.*, 177: 251–258.

- Kowalewska I, L Bielawski, and I Falandysz, 2007. Some elements and their bioconcentration factors in red aspen bolete *Leccinum rufum* from Northern Poland. *Bromat Chem. Toksykol*, 40: 329–335.
- Lynch SR and RD Baynes, 1996. Deliberations and evaluations of the approaches, endpoints and paradigms for iron dietary recommendations. *J. Nutrition*, 126: 2404–2409.
- Malinowska E, P Szefer and J Falandysz, 2004. Metals bioaccumulation by bay bolete, *Xerocomus badius*, from selected sites in Poland. *Food Chem.*, 84: 405–416.
- Manzi P, A Aguzzi and L Pizzoferrato, 2001. Nutritional value of mushrooms widely consumed in Italy. *Food Chem.*, 73: 321–325.
- Mendil D, ÖD Uluözlü, E Hasdemir and A Cağlar, 2004. Determination of trace elements on some wild edible mushroom samples from Kastamonu, Turkey. *Food Chem.*, 88: 281–285.
- Mendil D, ÖD Uluözlü, M Tüzen, E Hasdemir and H Sarı, 2005. Trace metal levels in mushroom samples from Ordu, Turkey. *Food Chem.*, 91: 463–467.
- Ouzouni PK, D Petridis, W Koller and KA Riganakos, 2009. Nutritional value and metal content of wild edible mushrooms collected from West Macedonia and Epirus, Greece. *Food Chem.*, 115:1575–1580.
- Pathak VN, N Yadav and M Gour, 1998. Mushroom production and processing tech. *Agrobotanica*, pp.2-3.
- Radulescu C, C Stihă, G Busuioc, AI Gheboianu, and IV Popescu, 2010. Studies concerning heavy metals bioaccumulation of wild edible mushrooms from industrial area by using spectrometric techniques. *Bull. Environ. Contam. Toxicol.*, 84: 641–646.
- Sesli E and M Tüzen, 1999. Levels of trace elements in the fruiting bodies of macrofungi growing in the East Black sea region of Turkey. *Food Chem.*, 65: 453–460.
- Sesli E, M Tüzen and M Soylak, 2008. Evaluation of trace metal contents of some wild edible mushrooms from Black sea region, Turkey. *J. Hazard. Mater.*, 160: 462–467.
- Shelly NJ, SMR Amin, MM Nuruddin, KU Ahmed and J Khandakar, 2008. Comparative Study on the Nutritional Composition of *Pleurotus ostreatus* (PO2) and Some Strains of *Pleurotus cystidiosus*. *Bangladesh J. Mushroom*, 2: 89-94.
- Shin CK, CF Yee, LJ Shya and M Atong, 2007. Nutritional properties of some edible wild mushrooms in Sabah. *J. Appl. Polym. Sci.*, 7: 2216–2221.
- Sivrikaya H, L Bacak, A Saraçbaşı, I Toroğlu and H Eroğlu, 2002. Trace elements in *Pleurotus sajorçaju* cultivated on chemithermomechanical pulp for bio-bleaching. *Food Chem.*, 79: 173–176.
- Soylak M, S Saracoglu, M Tüzen and D Mendil, 2005. Determination of trace metals in mushroom samples from Kayseri, Turkey. *Food Chem.*, 92: 649–652.
- Svoboda L, K Zimmermannova and P Kaláč, 2000. Concentrations of mercury, cadmium, lead and copper in fruiting bodies of edible mushrooms in an emission area of a copper smelter and a mercury smelter. *Sci. Total Environ.*, 246: 61–67.
- Türkekul I, M Elmastas and M Tüzen, 2004. Determination of iron, copper, manganese, zinc, lead, and cadmium in mushroom samples from Tokat, Turkey. *Food Chem.*, 84: 389–392.
- Tüzen M, 2003. Determination of heavy metals in soil, mushroom and plant samples by atomic absorption spectrometry. *Microchem. J.*, 74: 289–297.
- Tuzen M, E Sesli and M Soylak, 2007. Trace element levels of mushroom species from East Black Sea region of Turkey. *Food Control*, 18: 806–810.
- Tüzen M, I Turkekul, E Hasdemir, D Mendil and H Sari, 2003. Atomic absorption spectrometric determination of trace metal contents of mushroom samples from Tokat, Turkey. *Anal. Lett.*, 36: 1401–1410.
- Tüzen M, M Ozdemir and A Demirbas, 1998. Study of heavy metals in some cultivated and uncultivated mushrooms of Turkish origin. *Food Chem.*, 6: 247–251.
- Vetter J, 2003. Data on sodium content of common edible mushrooms. *Food Chem.*, 81: 589–593.
- World Health Organization (1982). Evaluation of Certain Food Additives and Contaminants (Twenty-sixth Report of the Joint FAO/WHO Expert Committee on Food Additives). WHO Technical Report Series, No.683, Geneva.
- Yamaç M, D Yıldız and C Sarıkürkcü, 2007. Heavy metals in some edible mushrooms from the Central Anatolia, Turkey. *Food Chem.*, 103: 263–267.
- Yaman M and I Akdeniz, 2004. Sensitivity enhancement in flame atomic absorption spectrometry for determination of copper in human thyroid tissues. *Anal. Sci.*, 20: 1363–1366.
- Zhu F, L Qu, W Fan, M Qiao, H Hao and X Wang, 2011. Assessment of heavy metals in some wild edible mushrooms collected from Yunnan Province, China. *Environ. Monit. Assess.*, 179: 191-199.